

NEW STABILITY CONDITIONS FOR POSITIVE CONTINUOUS–DISCRETE 2D LINEAR SYSTEMS

TADEUSZ KACZOREK

Faculty of Electrical Engineering
Białystok Technical University, ul. Wiejska 45D, 15–351 Białystok, Poland
e-mail: kaczorek@isep.pw.edu.pl

New necessary and sufficient conditions for asymptotic stability of positive continuous-discrete 2D linear systems are established. Necessary conditions for the stability are also given. The stability tests are demonstrated on numerical examples.

Keywords: positive systems, 2D linear systems, continuous-discrete systems.

1. Introduction

In positive systems inputs, state variables and outputs take only nonnegative values. A variety of models having positive systems behavior can be found in engineering, management science, economics, social sciences, biology and medicine, etc. An overview of the state of the art in positive systems is given in the monographs of Farina and Rinaldi (2000) as well as Kaczorek (2002).

Positive continuous-discrete 2D linear systems were introduced by Kaczorek (1998) along with positive hybrid linear systems (Kaczorek, 2007) and positive fractional 2D hybrid systems (Kaczorek, 2008a). Various methods of solvability of 2D hybrid linear systems were discussed by Kaczorek *et al.* (2008), and the solution to singular 2D hybrids linear systems was derived by Sajewski (2009). The realization problem for positive 2D hybrid systems was addressed by Kaczorek (2008b). Some problems of dynamics and control of 2D hybrid systems were considered by Dymkov *et al.* (2004) and Gałkowski *et al.* (2003). The problems of stability and robust stability of 2D continuous-discrete linear systems were investigated by Bistriz (2003), Busłowicz (2010a; 2010b, 2011) and Xiao (2001a; 2001b; 2003). Recently, stability and robust stability of a general model and of a Roesser type model of scalar continuous-discrete linear systems were analyzed by Busłowicz (2010a; 2010b; 2011).

In this paper, new necessary and sufficient conditions for asymptotic stability of positive continuous-discrete 2D linear systems will be presented.

The following notation will be used: \mathbb{R} is the set of real numbers, \mathbb{Z}_+ stands for the set of nonnegative inte-

gers, $\mathbb{R}^{n \times m}$ denotes the set of $n \times m$ real matrices, $\mathbb{R}_+^{n \times m}$ is the set of $n \times m$ matrices with nonnegative entries and $\mathbb{R}_+^n = \mathbb{R}_+^{n \times 1}$, I_n denotes the $n \times n$ identity matrix.

2. Preliminaries

Consider the linear autonomous continuous-discrete 2D system (Kaczorek, 1998; 2002)

$$\dot{x}(t, i+1) = A_0 x(t, i) + A_1 \dot{x}(t, i) + A_2 x(t, i+1), \\ t \in \mathbb{R}_+, \quad i \in \mathbb{Z}_+, \quad (1)$$

where $\dot{x}(t, i) = \partial x(t, i) / \partial t$, $x(t, i) \in \mathbb{R}^n$, $A_k \in \mathbb{R}^{n \times n}$ for $k = 0, 1, 2$.

Definition 1. The linear continuous-discrete 2D system (1) is called (*internally*) *positive* if $x(t, i) \in \mathbb{R}_+^n$, $t \in \mathbb{R}_+$, $i \in \mathbb{Z}_+$ for all initial conditions

$$x(0, i) \in \mathbb{R}_+^n, \quad i \in \mathbb{Z}_+, \\ x(t, 0) \in \mathbb{R}_+^n, \quad \dot{x}(t, 0) \in \mathbb{R}_+^n, \quad t \in \mathbb{R}_+. \quad (2)$$

Theorem 1. (Kaczorek, 1998; 2002) *The linear continuous-discrete 2D system (1) is positive if and only if*

$$A_2 \in M_n, \quad A_0, A_1 \in \mathbb{R}_+^{n \times n}, \\ A_0 + A_1 A_2 \in \mathbb{R}_+^{n \times n}, \quad (3)$$

where M_n is the set of $n \times n$ Metzler matrices (with non-negative off-diagonal entries).




Fig. 1. Shifting the zeros w into the unit circle of the complex plane.

The system (1) is called asymptotically stable if

$$\lim_{t \rightarrow \infty, i \rightarrow \infty} x(t, i) = 0.$$

Theorem 2. (Kaczorek, 2002) *The linear continuous-discrete 2D system (1) is asymptotically stable if and only if the zeros of the polynomial*

$$\begin{aligned} \det[I_n s z - A_0 - A_1 s - A_2 z] \\ = s^n z^n + a_{n,n-1} s^n z^{n-1} + a_{n-1,n} s^{n-1} z^n \\ + \dots + a_{10} s + a_{01} z + a_{00} \end{aligned} \quad (4)$$

are located in the left half of the complex plane s and in the unit circle of the complex plane z .

Theorem 3. (Kaczorek, 2002) *The positive linear system*

$$\dot{x} = Ax, \quad A \in M_n \quad (5)$$

is asymptotically stable if and only if the characteristic polynomial

$$\det[I_n s - A] = s^n + a_{n-1} s^{n-1} + \dots + a_1 s + a_0 \quad (6)$$

has positive coefficients, i.e., $a_k > 0$ for $k = 0, 1, \dots, n - 1$.

Lemma 1. (Farina and Rinaldi, 2000) *A nonnegative matrix $A \in \mathbb{R}_+^{n \times n}$ is asymptotically stable (a nonnegative Schur matrix) if and only if the Metzler matrix $A - I_n$ is asymptotically stable (a Metzler Hurwitz matrix).*

3. Main result

Theorem 4. *The positive linear continuous-discrete 2D system (1) is asymptotically stable if and only if all coefficients of the polynomial*

$$\begin{aligned} \det[I_n s(z+1) - A_0 - A_1 s - A_2(z+1)] \\ = s^n z^n + \bar{a}_{n,n-1} s^n z^{n-1} + \bar{a}_{n-1,n} s^{n-1} z^n \\ + \dots + \bar{a}_{10} s + \bar{a}_{01} z + \bar{a}_{00} \end{aligned} \quad (7)$$

are positive, i.e.,

$$\bar{a}_{k,l} > 0 \quad \text{for } k, l = 0, 1, \dots, n \quad (\bar{a}_{n,n} = 1). \quad (8)$$

Proof. It is well known that the zeros w_1, \dots, w_n of the characteristic polynomial

$$\det[I_n w - A] = w^n + a_{n-1} w^{n-1} + \dots + a_1 w + a_0 \quad (9)$$

located in the unit circle in the left half of the complex plane w can be shifted into the unit circle of the complex plane z by the substitution $w = z + 1$ (Fig. 1), i.e., the zeros z_1, \dots, z_n ($z_k = w_k + 1, k = 1, \dots, n$) of the characteristic polynomial

$$\begin{aligned} \det[I_n(z+1) - A] \\ = z^n + \hat{a}_{n-1} z^{n-1} + \dots + \hat{a}_1 z + \hat{a}_0. \end{aligned} \quad (10)$$

are located in the unit circle of the complex plane.

Note that the polynomial (7) is the characteristic polynomial of the positive system

$$\begin{aligned} \dot{x}(t, i+1) \\ = (A_0 + A_2)x(t, i) + (A_1 - I_n)\dot{x}(t, i) \\ + A_2 x(t, i+1). \end{aligned}$$

and its matrices $(A_0 + A_2), (A_1 - I_n), A_2$ are Metzger matrices. The sum of those matrices is also a Metzler matrix. Therefore, by Theorem 3 and the results of Kaczorek (2009), the positive continuous-discrete 2D system (1) is asymptotically stable if and only if the coefficients of the polynomial (7) are positive. ■

Example 1. Consider the system (1) with the matrices

$$\begin{aligned} A_0 &= \begin{bmatrix} 0.2 & 0 \\ 0.1 & 0.1 \end{bmatrix}, \\ A_1 &= \begin{bmatrix} 0.4 & 0 \\ 0.5 & 0.3 \end{bmatrix}, \\ A_2 &= \begin{bmatrix} -0.3 & 0 \\ 1 & -0.2 \end{bmatrix}. \end{aligned} \quad (11)$$

The matrices (11) satisfy the conditions (3) since

$$A_0 + A_1A_2 = \begin{bmatrix} 0.08 & 0 \\ 0.25 & 0.04 \end{bmatrix} \in \mathbb{R}_+^{2 \times 2}, \quad (12)$$

and then the system is positive.

In this case, the polynomial (7) has the form

$$\begin{aligned} & \det[I_n s(z+1) - A_0 - A_1 s - A_2(z+1)] \\ &= \det \begin{bmatrix} s(z+1) - 0.2 - 0.4s + 0.3(z+1) & \\ -0.1 - 0.5s - (z+1) & \\ & 0 \\ & s(z+1) - 0.1 - 0.3s + 0.2(z+1) \end{bmatrix} \quad (13) \\ &= s^2 z^2 + 1.3s^2 z + 0.5s z^2 + 0.42s^2 + 0.06z^2 \\ & \quad + 0.53s z + 0.13s + 0.05z + 0.01. \end{aligned}$$

All coefficients of the polynomial (13) are positive. Therefore, by Theorem 4, the positive continuous-discrete system (1) with (11) is asymptotically stable.

Theorem 5. *The positive continuous-discrete 2D linear system (1) is unstable if one of the following conditions is satisfied:*

- (i) $\det[-(A_0 + A_2)] \leq 0$,
- (ii) $\det[-A_2] \leq 0$,
- (iii) $\det[I_n - A_1] \leq 0$.

Proof. Substitution of $s = z = 0$ into (7) yields

$$\det[-(A_0 + A_2)] = \bar{a}_{00}. \quad (14)$$

If the condition (i) is satisfied, then from (14) we have $\bar{a}_{00} \leq 0$, and by Theorem 4 the system (1) is unstable. Substituting $s = 0$ into (7) we obtain

$$\begin{aligned} & \det[-A_2 z - (A_0 + A_2)] \\ &= \bar{a}_{0,n} z^n + \dots + \bar{a}_{01} z + \bar{a}_{00}, \quad (15) \end{aligned}$$

and $\det[-A_2] = \bar{a}_{0,n}$. If the condition (ii) is met, then $\bar{a}_{0,n} \leq 0$, and by Theorem 4 the system (1) is unstable. Similarly, substituting $z = 0$ into (7) we obtain

$$\begin{aligned} & \det[(I_n - A_1)s - (A_0 + A_2)] \\ &= \bar{a}_{n,0} s^n + \dots + \bar{a}_{10} s + \bar{a}_{00} \quad (16) \end{aligned}$$

and $\det[(I_n - A_1)] = \bar{a}_{n,0}$. If the condition (iii) is met then $\bar{a}_{n,0} \leq 0$ and, by Theorem 4, the system (1) is unstable. ■

Example 2. Consider the system (1) with the matrices

$$\begin{aligned} A_0 &= \begin{bmatrix} 0.5 & 0.3 \\ 0.4 & 0.4 \end{bmatrix}, \\ A_1 &= \begin{bmatrix} 0.2 & 0.1 \\ 0.1 & 0.3 \end{bmatrix}, \\ A_2 &= \begin{bmatrix} -0.3 & 0.1 \\ 0.2 & -0.4 \end{bmatrix}. \end{aligned} \quad (17)$$

The matrices (17) satisfy the conditions (3) since

$$A_0 + A_1A_2 = \begin{bmatrix} 0.46 & 0.28 \\ 0.43 & 0.29 \end{bmatrix} \in \mathbb{R}_+^{2 \times 2}, \quad (18)$$

and then the system is positive.

Using (17), we obtain

$$\begin{aligned} \det[-(A_0 + A_2)] &= \det \begin{bmatrix} -0.2 & -0.4 \\ -0.6 & 0 \end{bmatrix} = -0.24, \\ \det[-A_2] &= \det \begin{bmatrix} 0.3 & -0.1 \\ -0.2 & 0.4 \end{bmatrix} = 0.1, \\ \det[I_n - A_1] &= \det \begin{bmatrix} 0.8 & -0.1 \\ -0.1 & 0.7 \end{bmatrix} = 0.55, \end{aligned}$$

and the condition (i) of Theorem 5 is satisfied. Therefore, the positive system (1) with (17) is unstable.

In this case the polynomial (7) has the form

$$\begin{aligned} & \det[I_n s(z+1) - A_0 - A_1 s - A_2(z+1)] \\ &= \det \begin{bmatrix} sz + 0.8s + 0.3z - 0.2 & -0.1s - 0.1z - 0.4 \\ -0.1s - 0.2z - 0.6 & sz + 0.7s + 0.4z \end{bmatrix} \\ &= s^2 z^2 + 1.5s^2 z + 0.7s z^2 + 0.55s^2 + 0.1z^2 + 0.3s z \\ & \quad - 0.24s - 0.22z - 0.24, \end{aligned} \quad (19)$$

and, by Theorem 4, the system is also unstable.

4. Concluding remarks

New necessary and sufficient conditions for the asymptotic stability of continuous-discrete 2D linear systems have been established (Theorem 4). Some necessary conditions for asymptotic stability have also been given. The effectiveness of the new stability tests have been demonstrated on numerical examples. The deliberations can be also extended to fractional positive 2D continuous-discrete linear systems.

Acknowledgment

The author wishes to thank Professor M. Buśłowicz for his valuable comments and suggestions.

This work was supported by the Ministry of Science and Higher Education in Poland under the grant no. S/WE/1/11.

References

- Bitriz, Y. (2003). A stability test for continuous-discrete bivariate polynomials, *Proceedings of the International Symposium on Circuits and Systems*, Vol. 3, pp. 682–685.
- Buśłowicz, M. (2010a). Stability and robust stability conditions for a general model of scalar continuous-discrete linear systems, *Pomiary, Automatyka, Kontrola* **56**(2): 133–135.

- Busłowicz, M. (2010b). Robust stability of the new general 2D model of a class of continuous-discrete linear systems, *Bulletin of the Polish Academy of Sciences: Technical Sciences* **58**(4): 561–566.
- Busłowicz, M. (2011). Improved stability and robust stability conditions for a general model of scalar continuous-discrete linear systems, *Pomiary, Automatyka, Kontrola* **57**(2): 188–189.
- Dymkov, M., Gaishun, I., Rogers, E., Gałkowski, K. and Owens, D.H. (2004). Control theory for a class of 2D continuous-discrete linear systems, *International Journal of Control* **77** (9): 847–860.
- Farina, L. and Rinaldi, S. (2000). *Positive Linear Systems: Theory and Applications*, J. Wiley, New York, NY.
- Gałkowski, K., Rogers, E., Paszke, W. and Owens, D.H. (2003). Linear repetitive process control theory applied to a physical example, *International Journal of Applied Mathematics and Computer Science* **13** (1): 87–99.
- Kaczorek, T. (1998). Reachability and minimum energy control of positive 2D continuous-discrete systems, *Bulletin of the Polish Academy of Sciences: Technical Sciences* **46** (1): 85–93.
- Kaczorek, T. (2002). *Positive 1D and 2D Systems*, Springer-Verlag, London.
- Kaczorek, T. (2007). Positive 2D hybrid linear systems, *Bulletin of the Polish Academy of Sciences: Technical Sciences* **55**(4): 351–358.
- Kaczorek, T. (2008a). Positive fractional 2D hybrid linear systems, *Bulletin of the Polish Academy of Sciences: Technical Sciences* **56** (3): 273–277.
- Kaczorek, T. (2008b). Realization problem for positive 2D hybrid systems, *COMPEL* **27** (3): 613–623.
- Kaczorek, T. (2009). Stability of positive continuous-time linear systems with delays, *Bulletin of the Polish Academy of Sciences: Technical Sciences* **57**(4): 395–398.
- Kaczorek, T., Marchenko, V. and Sajewski, Ł. (2008). Solvability of 2D hybrid linear systems—Comparison of the different methods, *Acta Mechanica et Automatica* **2**(2): 59–66.
- Sajewski, Ł. (2009). Solution of 2D singular hybrid linear systems, *Kybernetes* **38** (7/8): 1079–1092.
- Xiao, Y. (2001a). Stability test for 2-D continuous-discrete systems, *Proceedings of the 40th IEEE Conference on Decision and Control, Orlando, FL, USA*, Vol. 4, pp. 3649–3654.
- Xiao, Y. (2001b). Robust Hurwitz–Schur stability conditions of polytopes of 2-D polynomials, *Proceedings of the 40th IEEE Conference on Decision and Control, Orlando, FL, USA*, Vol. 4, pp. 3643–3648.
- Xiao, Y. (2003). Stability, controllability and observability of 2-D continuous-discrete systems, *Proceedings of the International Symposium on Circuits and Systems, Bangkok, Thailand*, Vol. 4, pp. 468–471.



Tadeusz Kaczorek received the M.Sc., Ph.D. and D.Sc. degrees in electrical engineering from the Warsaw University of Technology in 1956, 1962 and 1964, respectively. In the years 1968–69 he was the dean of the Electrical Engineering Faculty, and in the period of 1970–73 he was a deputy rector of the Warsaw University of Technology. In 1971 he became a professor and in 1974 a full professor at the same university. Since 2003 he has been a professor at Białystok Technical University. In 1986 he was elected a corresponding member and in 1996 a full member of the Polish Academy of Sciences. In the years 1988–1991 he was the director of the Research Centre of the Polish Academy of Sciences in Rome. In 2004 he was elected an honorary member of the Hungarian Academy of Sciences. He has been granted honorary doctorates by several universities. His research interests cover the theory of systems and automatic control systems theory, especially singular multidimensional systems, positive multidimensional systems, and singular positive 1D and 2D systems. He initiated research in the field of singular 2D and positive 2D systems. He has published 24 books (six in English) and over 950 scientific papers. He has also supervised 69 Ph.D. theses. He is the editor-in-chief of the *Bulletin of the Polish Academy of Sciences: Technical Sciences* and a member of editorial boards of ten international journals.

Received: 24 January 2011

Revised: 9 March 2011